



Mehsana Unit :- 251, GIDC Estate, Mehsana-2, Gujarat, INDIA. Chhatral Unit :- B/23,24, Dharti Apollo Eco Industrial Park, Chhatral Kadi Road, Chhatral, Gujarat, INDIA.

Ph.: +91 9687335566, +91 9687335577 E-mail: info@morvellab.com • Web: www.morvellab.com









ISO 9001:2008 CERTIFIED COMPANY

INJECTION ANTIBIOTICS / ANTIMICROBIALS

OXYMOR Vet

Each ml. contains: Oxytetracycline 50 mg.

INDICATIONS: OXYMOR-Vet is one of the most potent broad spectrum antibiotics, effective against wide range of Gram positive and Gram negative bacteria. By parenteral route, peak plasma level is reached within one hour. It is useful in H.S, B. Q, Joint ill and other general infections.

ADMINISTRATION & DOSAGE: By I/M, I/V Use

Large Animal: 1 ml. per 10 Kg. body Wt. Small Animal: 1 ml. per 8 Kg. body Wt.



Available In: 100 ml, 30 ml.

OXYMOR LA

Each ml contains: Oxytetracycline Dihydrate 200 mg.

INDICATIONS: Oxymor LA is used in the treatment of infections in Catlle, Sheep, Goats and Pigs. Indicated in Contagious Bacterial infections like Hemorrhagic Septicemia, Black Quarter and Joint ill in sheep, goats and pigs. Also in Pneumonia and other general infections. When repeated treatment is not feasible, Oxymor LA is the drug of choice as an antibiotic.

ADMINISTRATION & DOSAGE: Use only as directed. Do not inject intravenously. Oxymor LA is designed for intramuscular injection. Normally, a single treatment of 1 ml/10 kg body mass will be sufficient for the treatment of susceptible disease conditions. Repeat the dose after 72 hours.



Available In : 50 ml.

GENTAMOR® Vet

Each ml. contains: Gentamicin 40 mg.

INDICATIONS: GENTAMOR-Vet is an excellent medicine against Gram Negative Bacteria. A bactericidal antibiotic active even at low concentrations against wide spectrum of bacteria. Gentamor Vet. is the best antibiotic against infections caused by bacteria in U.T., R.T. It is the drug of choice to treat Metritis & Repeat Breeders.

ADMINISTRATION & DOSAGE: By I.M. & I.V. Use only Large Animal: 1-2 mg./ Kg. body Wt. Small Animal: 2-4 mg./Kg. body Wt.

Intra-Uterine: 2ml. Gentamor-Vet diluted with 20ml of Distilled water in case of repeat breeders.



Available In: 100 ml, 30 ml.

KANCIN Vet

Each ml contains: Amikacin Sulphate I.P: 250 mg.

Indications: KANCIN Vet is indicated in Gynaecological infections such as: METRITIS, CERVICITS, VAGINITIS. It is also effective in Urinary Infections. It is useful in MASTITIS, ENTERITIS, Osteomyelitis etc. Highly effective against Gentamicin resistant organisms.

Administration & Dosage: For I.M & I.V Use only. Large Animals: 5-7 mg/ Kg Body Wt./ 12 hr. Single Dose: 10 – 15 ml Small Animals: 10 mg/ Kg Body Wt / 12 hr



Available In : 100 ml.

MENDRO Vet

Each ml. contains: Enrofloxacin 100 mg.

INDICATIONS: MENDRO-Vet is a broad spectrum, bactericidal and mycoplasmicidal antibiotic. It is highly effective in any Urinogenital & Respiratory Tract Infections. It is indicated in Pneumonia, Metritis, Endometritis, Cystitis and Mastitis. It is also useful in any general bacterial & Mycoplasmal Infections.

ADMINISTRATION & DOSAGE: By I.M. Use only. Large Animal: 2.5-5 mg/Kg B.Wt (1ml/ 20 kg. body Wt.)



Available In : 100 ml.

NEOSPA

Each ml. contains: Neomycin 140 mg.

INDICATIONS: NEOSPA is a most stable antibiotic to treat acute & chronic bacterial infections due to organisms susceptible to Neomycin. It is useful in Pneumonia, Bronchitis, Pleurisy and bacterial complications associated with viral diseases as well as Mastitis & Metritis.

ADMINISTRATION & DOSAGE: By I.M & I.V use. **Large Animal**: 10-20 mg/Kg. B.Wt. **Small Animal**: 20-40 mg/

Kg. B.Wt.



Available In: 30 ml.

CEFVEL Vet

Each vial contains : Cefoperazone Sodium 3gm+ Sulbactum Sodium 1.5gm

INDICATIONS: CEFVEL with the combination of Cefoperazone & Sulbactum exhibits a marked synergic action. Cefoperazone is a semi synthetic broad spectrum Cephalosporin antibiotic. Sulbactum Sodium nullifies the action of any resistance by betalactamase. CEFVEL successfully treats Mastitis, Metritis, Cervicitis, Abdominal infection, Pneumonia and septicaemia.

ADMINISTRATION & DOSAGE: By I/M., I.V. use only. **Large Animal**: 5-10 mg./kg. body wt.

Small Animal: 10-15 mg/kg body wt.

Dissolve the content with sterile water provided along with the pack & use.

Available In : 4.5 gm, 3 gm.



MORCEF Vet

Each vial contains: Ceftriaxone Sodium I.P: 4000 mg.

Indications: Ceftriaxone is a third-generation antibiotic from the Cephalosporin family of antibiotics. Its main feature is long plasma half-life.

MORCEF Vet is highly indicated in Mastitis, Metritis, Pyometra, Septicemia, Broncho Pneumonia, Urinary Tract Infections, Osteo arthritis and Bacterial Meningitis.

Administration & Dosage:

By I/M, & I/V use only.

Large Animals: 5 - 10 mg / Kg. Body wt. (Approx. 4 Gm)

Small Animals: 15 – 25 mg/ Kg. Body wt.

Dissolve the content with sterile water provided along with the pack and use.



Available In : 4 gm.

INJECTION ANTIBIOTICS / ANTIMICROBIALS

LOXY Vet

Each vial contains : Amoxycillin 2.25 gm. + Cloxacillin 2.25 gm.

INDICATIONS: LOXY-Vet is very effective against both Gm+ve as well as Gm-ve Bacterial spectrum. It is indicated in MASTITIS, METRITIS, Salmonellosis, Calf scour, E. Coli infections and post surgical cases. It is the drug of choice for all Resp Tract infections as well as Uninary Tract infections.

ADMINISTRATION & DOSAGE: By I.M & I.V Use only.

Large Animal: 6-10 mg/kg. B.Wt. Small Animal: 10-20 mg/kg B.Wt.

Dissolve the content with sterile water for injection.



Available In : 4.5 gm.

NELLIN® Vet

Each vial contains: Ampicillin 2.25gm + Cloxacillin 2.25gm

INDICATIONS: This combination of 2 antibiotics is the drug of choice for Metritis, Retained Placenta, Mastitis and Postsurgical treatment. It is very effective in H.S, B.Q, Brucellosis, Resp. tract infections and Pyrexia due to unknown origin. It is active against both Gm +ve and Gm-ve Bacteria.

ADMINISTRATION & DOSAGE: By I.M & I.V Use only.

Large Animal: 6-10 mg/Kg.body wt. **Small Animal:** 10-20 mg/Kg body wt.

Dissolve the content with sterile water for injection.



Available In : 4.5 gm, 3 gm.

MORPRIM Vet

Each 5 ml contains: Inj. of Trimethoprim 80 mg + Sulphamethaxazole 400 mg.

INDICATIONS: MORPRIM-Vet is a combination of two bactericidal chemotherapeutics, Trimethoprim & Sulphamethoxazole. It is the drug of choice to treat Diarrhoea & H.S.

ADMINISTRATION & DOSAGE: By I/V use only. 15 – 30 mg/Kg.B.Wt (15 ml for 400 Kg animal)



Available In : 30 ml.

PROTONIL Vet RTU

Each ml contains : Diminazene Aceturate 70 mg. + Phenazone B.P. 375 mg.

INDICATION: Protonil Vet is an effective Anti haemoprotozoal, for the treatment and control of Babesiosis, Trypanosomiasis and Theilariosis in animals.

Protonil as the name suggests nullifies the causative protozoal organisms in the blood. Protonil has also antibacterial and antifungal actions. Phenazone in Protonil is an antipyretic and brings down the fever occurring due to protozoal infections.

ADMINISTRATION & DOSAGE: By deep I/M or S/C use. In horse: By Intramuscular only

LARGE ANIMAL: 1ml/10-20 Kg Body weight in Total dose should not exceed: 9 g/Animal/day.

Contra indicated in Camels.



Available In : 30 ml, 90 ml.

INJECTION ANALGESICS

MELOX Vet

Each ml. contains: Meloxicam B.P. 5 mg.

INDICATIONS: MELOX-Vet is a non-steroidal antiinflammatory drug, which selectively inhibits COX-2 enzyme and has no side effects. It is used to reduce the inflammatory muscular pain and fever. Along with antibiotic, it is useful for the treatment of PNEUMONIA, Bronchopneumonia, SPRAIN, ARTHRITIS, MASTITIS, Prolapse of Uterus and Otitis.

BW (30 ml/300 Kg BW)

IN OTHER CONDITIONS: 15 ml for 300 Kg BWt. It is to be repeated daily or as directed by the Veterinarian.

Avoid in late pregnancy.



Available In: 100 ml, 30 ml.

MELOX® PLUS

Each ml contains: Meloxicam 5 mg. + Paracetamol 150 mg+ Lignocaine: 1% w/v

INDICATIONS: MELOX PLUS is a non-steroidal antiinflammatory drug with Paracetamol combination and is used to reduce the infla-mmatory muscular pain and fever. Along with antibiotics, it is useful for the treatment of Pneumonia, Prolapse, Arthritis, Mastitis, Laminitis, Myositis and Otitis.

ADMINISTRATION & DOSAGE: By I/M only LARGE ANIMAL: 15-20 ml per animal (300 Kg B.Wt) SMALL ANIMAL: 0.5-2 ml per animal.

Metroisan and Paracetanid Injection
MEL

Available In: 100 ml, 30 ml.

MORGAN® Vet

Each ml. contains: Dicyclomine Hydrochloride 10 mg.

INDICATIONS: MORGAN-Vet injection has rapid & sustained antispasmodic effect on the smooth muscles. It is excellent for the treatment of painful spasmodic conditions of the G.I.Tract, Colic and other painful conditions.

ADMINISTRATION & DOSAGE : I.M use only.

LARGE ANIMAL: 0.5 mg/ Kg. BW (15 ml/300 Kg Cow, Buffalo)

SMALL ANIMAL: 1 to 2 ml.

MORGAN should not be mixed with other drugs in the same syringe.



Available In : 30 ml.

UDIL-S Vet

Each ml. contains: Phenyl Butazone 200 mg. + Sodium Salicylate 20 mg.

INDICATIONS: A unique combination to relieve the pain, fever and swelling. It is indicated in Myositis, Laminitis, Neuritis and other such inflammatory conditions. It is useful in the treatment of Non-visceral Trauma, Pain, Lameness, Swelling and Pyrexia.

ADMINISTRATION & DOSAGE: By deep I.M. use only. LARGE ANIMAL: 10-15 ml.

SMALL ANIMAL: 1-2 ml.



Available In: 100 ml, 30 ml.

MORPIN Vet

Each ml. Contains: Analgin 500 mg

INDICATIONS: MORPIN-Vet has central analgesic action. It increases blood supply to striped muscles which is therapeutically desirable. It is used for relief of pain in chronic arthritis, rheumatic conditions of muscles & joints.

ADMINISTRATION & DOSAGE: I/M use only

LARGE ANIMAL: 10 – 20 mg/ Kg. BW (10 -15 ml / 300 Kg [

Available In : 100 ml.

Cow, Buffalo) SMALL ANIMAL: 0.5 – 2 mg/Kg BW (0.5 - 2 ml-Total Dose)

INJECTION ANAESTHETICS

MORCAIN Vet

Each ml. contains : Lignocaine Hcl 21.33 mg. \pm

Adrenaline Bititrate 0.01 mg.

INDICATIONS: MORCAIN-Vet is a potent sedative. It is an agent for conduction of anesthesia in minor surgical procedures.

ADMINISTRATION & DOSAGE : S/C Use : The dosage depends on the indication & the area to be anaesthetised.



Available In: 30 ml.

SICVEL Vet

Each ml. contains: Triflupromazine HCl 20 mg.

INDICATIONS: SICVEL-Vet is used in the Veterinary field for the management of psycho motor over activity, usually during clinical procedures, pre-operative agent, combination with local or general anesthesia. It increases tolerance of animals to pain. Prolapse of Uterus or Vagina. ADMINISTRATION & DOSAGE: I/M, I/V. Use

LARGE ANIMAL: 0.22-0.33 mg/Kg BW. (2 – 3 ml only for 300 Kg BW)

SMALL ANÍMAL: 2-4 mg/Kg BW

(0.5 - 2 ml – Total Dose) Maximum dosage level may vary from animal to animal & is to be adjusted as per requirement.



Available In : 5 ml.

INJECTION VITAMINS

ADVEL-A

Each 2 ml. contains: Highly concentrated Vitamin A 600000 I.U.

INDICATIONS : ADVEL-A In Infertility and supportive treatment in R.T. GI and UT infections.

ADMINISTRATION & DOSAGE : I/M use only.

LARGE ANIMAL: In Infertility: 2 amp (4ml) on alternate days for 2 weeks In other conditions: 3 - 6 amp per week.



Available In : 6 amp x 2 ml.

ADVIT-DE+H

Each ml contains: Vitamin A 250,000 I.U + Vitamin D3 25,000 I.U + Vitamin E 100 I.U + Vitamin H (Biotin) 12.5 mg.

Indications: ADVIT-DE+H is an injection to promote Maintenance, Growth, Lactation, Reproduction and well being of animals. Vitamin A is essential for normal functioning of the epithelial cells. Vitamin D3 is essential for growth and absorption of Calcium Vitamin E is an antioxidant and helps in the treatment of Muscular Dystrophy Vitamin H is a nitrogen-containing acid essential for growth and well-being in animals. It also helps in gluconeogenesis. Sesame oil used for better animal compliance.

ADMINISTRATION & DOSAGE : For I.M Use only [Deep I/M route].

LARGE ANIMAL: 5ml /day for Prophylaxis and 10 ml for treatment

SMALL ANIMAL: 1-2 ml/day as Prophylaxis and 2-4 ml for treatment. To be given on alternate days for 2 weeks. It should be repeated after 4 weeks if needed. Contraindicated in Neonates.



Available In : 10 ml.

ADVIT-DE®

Each ml. contains : Vitamin A 2,50,000 I.U. \pm Vitamin D3 25,000 I.U. \pm

Vitamin E 100 I.U.

INDICATIONS: ADVIT-DE is specially indicated for maintaining the growth, production, and lactation of farm animals. It is useful for the treatment of under developed genitalia, infertility. It maintains the pregnancy. It is also useful as a supportive therapy in General Debility, Stress and as a co-therapy with antibiotics.

ADMINISTRATION & DOSAGE: By deep I/M Use only. LARGE ANIMAL: 5ml. per day for Prophylaxis.10 ml for treatment.

SMALL ANIMAL: 1-2 ml. / day as Prophylaxis 2-4 ml for treatment. To be given on alternate days for 2 weeks and should be repeated after 4 weeks if needed.



Available In : 10 ml.

BITEEX® Vet

Inj. of Vit B complex Each ml. contains: Thiamine HCl 33 mg. + Riboflavine Sod.Phos. 1.66 mg. + Pyridoxine HCl 3.3 mg. + Nicotinamide 33.3 mg. + Cyanocobalamine 33.3 mg. + D-Panthenol 1.66 mg.

INDICATIONS: BITEEX is a high potency Vitamin B Complex Being IV compatable, it ensures a quick action. It is useful as a Co-therapy with antibiotics for recovery for infections. It is indicated in Anorexia, Anaemia & General Debility. It is also useful in treating diseases due to B Complex deficiency and as a Vitamin supplement.

ADMINISTRATION & DOSAGE:

I/V & I/M Use Only.

LARGE ANIMAL: 15 - 20 ml. SMALL ANIMAL: 2-5 ml.



Available In: 100 ml, 30 ml.

BELIVE Vet

Inj of Crude liver extract with Vitamin B Complex.

Each ml contains: Vit B1 (Thiamine Hcl): 10 mg + Vit B2 (Riboflavin): 3 mg + Vit B3 (Niacinamide) 100 mg + Vit B12 (Cyanocobalamin: 10 mcg + Liver Injection Crude: 0.66 ml (eq to 2 mcg of Vit B12 per ml)

INDICATIONS: BELIVE Vet is an injection of Crude liver extract with vitamin B-complex. It is specially for the treatment of anorexia and liver disorders associated with fatty degeneration of liver, sluggish liver. debility and general Weaknesses & Blood Protozoan diseases.

ADMINISTRATION & DOSAGE: Deep I/M Use only.

LARGE ANIMAL: 5 to 10 ml. Daily SMALL ANIMAL: 1 to 2 ml. Daily



Available In : 100 ml, 30 ml.

MORBIC-C

Each ml contains: Ascorbic acid 250 mg

INDICATION: MORBIC-C is an anti oxidant. It enhances tissue healing and growth. Improves immune function and acts as an anti-inflammatory agent. Supportive treatment during all treatments.

ADMINISTRATION & DOSAGE: By I/M use only LARGE ANIMALS: Give 20 – 30 ml twice weekly. SMALL ANIMALS: Give 3 – 5 ml twice weekly



Available In : 30 ml.

INJECTION VITAMINS

BION-12 Plus

Each ml contains: Thiamine Hydrochloride (Vitamin B1) I.P 50 mg + Pyridoxine Hydrochloride (Vitamin B6) I.P. 50 mg + Cynocobalamine (Vitamin B12) I.P. 500 mcg + Benzyl Alcohol I.P. 1.5 % +

Water for Injection I.P. Q.S.

INDICATION: Bion 12-Plus is a combination of three vital vitamins namely Vitamin B1, B6 and B12. It is mainly indicated for neurological disorders, anorexia, fatigue and debility conditions. It is also indicated in, dermatitis, macrocytic anaemia, pregnancy and lactation. It is useful for Nervine Stability.

ADMINISTRATION & DOSAGE: For I.M. use only. LARGE ANIMALS: 10 ml I.M. Daily

SMALL ANIMALS: 2 to 3 ml I.M. Daily.



Available In: 100 ml.

INJECTION MINERALS

CALCIVET

Inj. of Calcium Each ml. Contains:

Calcium Levulinate 76.4 mg. + Vit D3 5000 I.U. + Vit B12 50 mcg.

INDICATIONS: CALCIVET injection is a unique combination of calcium with Vit. D3 and Vit. B12 to fulfill the multiple needs of animals particularly lactating animals. During late pregnancy and early lactation, requirement of calcium is very high. CALCIVET is used for the prevention and treatment of hypocalcaemia in cows and buffaloes particularly during pre and post parturition period. As a tonic to maintain milk yield & to improve general health of the animal in conditions of debility & weakness.

ADMINISTRATION & DOSAGE: Deep I/M Use only.

LARGE ANIMAL: Pre Parturition: 5-10 ml.

Thrice a week. Post Parturition: 15-20 ml. thrice a week.

DEBILITATED ANIMAL: 10 ml. thrice a week. SMALL ANIMAL: 1-5 ml. thrice a week.



Available In: 100 ml, 30 ml.

TENOPHOS Vet

Each ml. contains: Sodium Acid Phosphate 40.3% w/v eg. to elemental Phosphate 80 mg. per ml.

INDICATIONS: TENOPHOS-Vet contains in-organic phosphorus, which is more useful in acute cases. It is indicated for Infertility cases such as: Under-developed Genitalia, Delayed maturity, Anoestrus and maintenance of Pregnancy. It is very effective to treat Metabolic disorders such as: Post Parturient Haemoglobinuria, Hypo Phosphataemia, Ketosis & Milk Fever. It is useful in General debility conditions and in the treatment of Bloat, Tetany, Pica & Rickets in lambs.



LARGE ANIMAL: 5-10 ml daily for alternate days.

SMALL ANIMAL: 2-5 ml daily Dose can be adjusted according to the phosphorus need of the animal. It is advisable to administer half the dose by I/V and remaining half dose by s/c route in several places.

INJECTION OTHERS

MORDEX Vet

Each ml. Contains: Dexamethasone 4 mg.

INDICATIONS : MORDEX-Vet is a highly active glucocorticoid with powerful anti- inflammatory and anti-ketotic (gluconeogenesis) effect. It is useful in ketosis, inflammation, surgical shock and pregnancy toxaemia.

ADMINISTRATION & DOSAGE: I/M., slow I/V Use. LARGE ANIMAL: Total Dose: 10 - 20 mg (2-5 ml) SMALL ANIMAL: Total Dose: 0.5-2 mg (0.5 to 2 ml),



Available In : 30 ml, 5 ml.

ALERT Vet

Each ml. Contains: Pheniramine Maleate 22.95 mg.

INDICATIONS: Of the many antihistaminic preparations, ALERT Vet, is the best for its mighty, rapid and prolonged action with excellent tolerance. It is used for allergic reactions like Itching of unknown genesis. Dermatitis, Skin Oedema, Insect bites, Bloat in ruminants due to histamine, atony of the rumen and also Mastitis

ADMINISTRATION & DOSAGE: I/M I/V Use

LARGE ANIMAL: 5-10 ml. SMALL ANIMAL: 0.5-2 ml. Available In: 100 ml, 30 ml.



RELIN Vet

Each ml contains: Buserelin acetate eq. to Buserelin 4 mcg.

Indications: Relin is basically a Gonadotropin Releasing Hormone Analogue (GnRH). It is used to induce heat in dairy cows and buffaloes. It is indicated to improve the conception rate (at the time of AI), to treat Ovarian cyst, delayed ovulation and anovulation.

Administration & Dosage: For I.M., I.V. and S.C. use only. To improve conception rate after A.I: 5ml In the case of ovarian cyst related irregulations: 2.5 ml Delayed Ovulation, Anovulation: 2.5 ml



Available In : 5 ml.

PULP DEPOT

Each ml. contains: Hydroxy Progesterone Caproate 250 mg.

INDICATIONS: A real approach to solve Reproductive Problems like Habitual abortion, Repeat breeders, prolapse of uterus, induction of Oestrus. Infertility caused by Progesterone deficiency.

ADMINISTRATION & DOSAGE: For Large Animal: I/M route only. Habitual Abortion in early pregnancy: 750 mg. PULP DEPOT I/M after 1½ month of pregnancy to be repeated 4 or 5 times at 10 days interval. Repeat breeders with weak corpus Luteum (Luteal Defeciency): 500 mg. or 750 mg. PULP DEPOT I/M to be followed for 3 weeks. Prolapse of Uterus: 750 mg. for two days. Induction of Oestrus: 500 mg. PULP DEPOT. If not responded repeatafter 10 days. For Sheep and Goats: 100 mg-200 mg only (0.5 ml to 1 ml)





Available In : 3 ml, 2 ml, Amp.

INJECTION OTHERS

METTLE® Vet

Each ml. contains: Methyl Ergometrine Maleate 0.2 mg.

INDICATIONS: A highly valuable Ecbolic [Uterine stimulant] for Veterinary Obstetrics to stimulate uterus, hasten parturition and involution of uterus. To prevent excessive haemorrohage and Retention of Placenta.

ADMINISTRATION & DOSAGE: By I.M. / I.V. Management of Third stage of Labour: For Cow and Buffalo: 10-15ml. Preferably by I.V. For sub involution of Lochiometra & Puerperal Bleeding: 5-10 ml. Daily for 3-5 days by I.M.



Available In : 5 ml.

REMOT Vet

Each ml. contains : Frusemide 50 mg.

INDICATIONS: REMOT Vet is a diuretic for parenteral administration in the treatment of oedema of all types. Congestive Oedema, pathological Oedema of Udder and perineum of cows and mares. For supportive treatment in elimination of fluid from the body cavities.

ADMINISTRATION & DOSAGE: I/M, I/V Use LARGE ANIMAL: 1-2 mg/ KG BW (5-10 ml) SMALL ANIMAL: 2-4 mg / Kg BW. (1-3 ml)



Available In : 10 ml, 30 ml.

ATROPINE SULPHATE

Each ml. contains: Atropine Sulphate 1 mg.

INDICATIONS: As an antidote to organophos phorous poisoning and Co-therapy in diarrohoea by reducing peristalsis movement.

ADMINISTRATION & DOSAGE:

By I/M, Slow I/V use only. For the treatment of Diarrhea: LARGE ANIMAL: 30 - 40 ml For treatment of Poisoning:

LARGE ANIMAL: 0.01 - 0.1 mg / Kg Body Wt. SMALL ANIMALS: 0.02 - 0.04 mg / Kg Body Wt



Available In: 30 ml, 100 ml.

CHROME

Each ml. contains: Carbazochrome Salicylate eq. to Adrenochrome Monosemicarbazone I.P. 5 mg.

INDICATIONS: CHROME is a coagulant used in the treatment of Capillary Haemorrhages characterised by increased capillary permeability.

ADMINISTRATION & DOSAGE: By I/M use. LARGE ANIMAL: Total Dose: 20-25 mg

Normally 5-10 ml

SMALL ANIMAL: Total Dose: 5-10 mg



Available In: 10 ml, 30 ml.

CHROME PLUS

Each ml contains: Carbazochrome Salicylate eg. to Adrenochrome Monosemicarbazone I.P 10 mg.

Indications: Chrome Plus is a coagulant used to arrest capillarity haemorrhages. The advantage of Chrome plus is that it has the double strength of Carbazochrome Salicylate. It ensures quick action during bleeding in the surgical procedures.

Administration & amp; Dosage: For I.M. use only.

Large animals: 20 - 25 mg (Total dose) Small animals: 5 - 10 mg (Total dose)



Available In : 30 ml.

IVEL Vet

Each ml. contains: Ivermectin 10 mg.

INDICATIONS: IVEL-Vet is a ready to use Anthelmintic and endectocide that exhibits activity against both Endoparasites and Ectoparasites. It is indicated in G.I. round worms. Lungworms. Screw worms. It is effective against ectoparasites such as : Lice, Mites, Ticks and

ADMINISTRATION & DOSAGE: By S/C use only. IVEL Vet is not to be used Intravenously or

Intramuscularly.

LARGE & SMALL ANIMAL: 200mcg/Kg BW. (10mg / 50 KG BW) 1ml for 50 Kg BWt



Available In: 10 ml, 50 ml, 100 ml.

LESOL-75

Each ml. contains: Levamisole HCI 75 mg.

INDICATIONS: LESOL-75 is a broad spectrum anthelmintic used for the treatment and routine control of nematode infections. Provides greater resistance to diseases like FMD & R P by increasing cell mediated immunity and antibody titre.

ADMINISTRATION & DOSAGE: By S.C. route 1 ml/30 Kg. body wt. If req. repeat after 48 hours.

LARGE ANIMAL: 3 to 5 mg/Kg BW (10-20 ml for 300

Kg Animal)



Available In: 30 ml.

CONTACT US:

